Diagnóstico de la fase preclínica de la degeneración mixomatosa de la válvula mitral

Jorge Pérez Nácher

Veterinario con especial interés en cardiología clínica de pequeños animales

Bibliografía

1. Keene BW, Atkins CE, Bonagura JD, Fox PR, Häggström J, Luis Fuentes V, et al. ACVIM consensus guidelines for the diagnosis and treatment of myxomatous mitral valve disease in dogs. J Vet Intern Med. 2019;33(3):1127-1140. doi:10.1111/jvim.15488.

2. Boswood A, Häggström J, Gordon SG, Wess G, Stepien RL, Oyama MA, et al. Effect of pimobendan in dogs with preclinical myxomatous mitral valve disease and cardiomegaly: the EPIC study-a randomized clinical trial. J Vet Intern Med. 2016;30(6):1765-1779. doi:10.1111/jvim.14586.

3. Ljungvall I, Rishniw M, Porciello F, Ferasin L, Ohad DG. Murmur intensity in small-breed dogs with myxomatous mitral valve disease reflects disease severity. J Small Anim Pract. 2014;55(11):545-550. doi:10.1111/jsap.12265.

4. Birkegård AC, Reimann MJ, Martinussen T, Häggström J, Pedersen HD, Olsen LH. Breeding restrictions decrease the prevalence of myxomatous mitral valve disease in Cavalier King Charles Spaniels over an 8- to 10-year period. J Vet Intern Med. 2016;30(1):63-68. doi:10.1111/jvim.13663.

5. Lord PF, Hansson K, Carnabuci C, Kvart C, Häggström J. Radiographic heart size and its rate of increase as tests for onset of congestive heart failure in Cavalier King Charles Spaniels with mitral valve regurgitation. J Vet Intern Med. 2011;25(6):1312-1319. doi:10.1111/j.1939-1676.2011.00792.x.

6. Malcolm EL, Visser LC, Phillips KL, Johnson LR. Diagnostic value of vertebral left atrial size as determined from thoracic radiographs for assessment of left atrial size in dogs with myxomatous mitral valve disease. J Am Vet Med Assoc. 2018;253(8):1038-1045. doi:10.2460/javma.253.8.1038.

7. Stepien RL, Rak MB, Blume LM. Use of radiographic measurements to diagnose stage B2 preclinical myxomatous mitral valve disease in dogs. J Am Vet Med Assoc. 2020;256(10):1129-1136. doi:10.2460/javma.256.10.1129.

8. Jang IS, Yoon WK, Choi EW. N-terminal pro-B-type natriuretic peptide levels in normotensive and hypertensive dogs with myxomatous mitral valve disease stage B. Ir Vet J. 2023;76(1):3. doi:10.1186/s13620-023-00233-0.

9. Hansson K, Häggström J, Kvart C, Lord P. Left atrial to aortic root indices using two-dimensional and M-mode echocardiography in Cavalier King Charles Spaniels with and without left atrial enlargement. Vet Radiol Ultrasound. 2002;43(6):568-575. doi:10.1111/j.1740-8261.2002.tb01051.x.

10. Cornell CC, Kittleson MD, Della Torre P, Häggström J, Lombard CW, Pedersen HD, et al. Allometric scaling of M-mode cardiac measurements in normal adult dogs. J Vet Intern Med. 2004;18(3):311-321. doi:10.1892/0891-6640(2004)18<311:ASOMCM>2.0.CO;2.

11. Reynolds CA, Brown DC, Rush JE, Fox PR, Nguyenba TP, Lehmkuhl LB, et al. Prediction of first onset of congestive heart failure in dogs with degenerative mitral valve disease: the PREDICT cohort study. J Vet Cardiol. 2012;14(1):193-202. doi:10.1016/j.jvc.2012.01.008.

12. Borgarelli M, Ferasin L, Lamb K, Chiavegato D, Bussadori C, D'Agnolo G, et al. The predictive value of clinical, radiographic, echocardiographic variables and cardiac biomarkers for assessing risk of the onset of heart failure or cardiac death in dogs with preclinical myxomatous mitral valve disease enrolled in the DELAY study. J Vet Cardiol. 2021;36:77-88. doi:10.1016/j.jvc.2021.04.009.

13. Hori Y, Iguchi M, Hirakawa A, Kamiya Z, Yamano S, Ibaragi T, et al. Evaluation of atrial natriuretic peptide and cardiac troponin I concentrations for assessment of disease severity in dogs with naturally occurring mitral valve disease. J Am Vet Med Assoc. 2020;256(3):340-348. doi:10.2460/javma.256.3.340.